

# Condensed Matter Physics

## Assignment I

Due: November 25, 2020

### 1. Bose-Einstein Distribution

Derive the occupation probability of energy levels for a collection of bosons

$$f(E) = \frac{1}{\exp\left(\frac{E-\mu}{k_B T}\right) - 1}, \quad (1)$$

where  $T$  is the temperature,  $\mu$  is the chemical potential and  $k_B$  is the Boltzmann constant.

(10 Marks)

### 2. Poisson Distribution

In the Drude model the probability of an electron suffering a collision in any infinitesimal interval  $dt$  is  $dt/\tau$ .

- Show that an electron picked at random at a given moment had no collision during the preceding  $t$  seconds with probability  $\exp(-t/\tau)$ . Show that it will have no collision during the next  $t$  seconds with the same probability.
- Show that the probability that the time interval between two successive collisions of an electron falls in the range between  $t$  and  $t + dt$  is  $(dt/\tau) \exp(-t/\tau)$ .
- Show as a consequence of (a) that at any moment the mean time back to the last collision (or up to the next collision) averaged over all electrons is  $\tau$ .
- Show as a consequence of (b) that the mean time between successive collisions of an electron is  $\tau$ .
- Derive the probability distribution for  $T$ , the time between the last and next collision.

(15 Marks)

### 3. Joule Heating

Consider a metal at uniform temperature in a static uniform electric field  $E$ . An electron experiences a collision, and then, after a time  $t$ , a second collision. In the Drude model, energy is not conserved in collisions, for the mean speed of an electron emerging from a collision does not depend on the energy that the electron acquired from the field since the time of the preceding collision.

- Show that the average energy lost to the ions in the second of two collisions separated by a time  $t$  is  $(eEt)^2/2m$  (the average is over all directions in which the electron emerged from the first collision).
- Show that the average energy loss to the ions per electron per collision is  $(eEt)^2/m$ , and hence that the average loss per cubic centimeter per second is  $(ne^2\tau/m)E^2 = \sigma E^2$ . Deduce that the power loss in a wire of length  $L$  and cross section  $A$  is  $I^2 R$ , where  $I$  is the current flowing and  $R$  is the resistance of the wire.

(15 Marks)